arXiv:1401.6544v1 [math.AG] 25 Jan 2014

BIFURCATION LOCUS AND BRANCHES AT INFINITY OF A POLYNOMIAL $f : \mathbb{C}^2 \to \mathbb{C}$

ZBIGNIEW JELONEK AND MIHAI TIBĂR

Abstract. We show that the number of bifurcation points at infinity of a polynomial function $f : \mathbb{C}^2 \to \mathbb{C}$ is at most the number of branches at infinity of a generic fiber of f and that this upper bound can be diminished by one in certain cases.

1. Introduction

Let $f : \mathbb{C}^2 \to \mathbb{C}$ be a polynomial function in a fixed coordinate system. It is well known (as being proved originally by Thom [T]), that f is a locally trivial C^∞ fibration outside a finite subset of the target. The smallest such set is called *the bifurcation set of* f and will be denoted here by $B(f)$. The set $B(f)$ might be larger than the set $f(\mathrm{Sing}f)$ of critical values of f since it contains also the set $B_\infty(f)$ of *bifurcation points at infinity*. Roughly speaking, $B_\infty(f)$ consists of points at which the restriction of f to a neighbourhood of infinity (i.e. outside a large enough ball) is not a locally trivial bundle. We say that $a \in B_\infty(f)$ is a *critical value at infinity* of f. There are several criteria to detect such a value, one may consult e.g. [Su], [HL], [Ti1], [Du], [CK], [Ti2]. For instance, $a \in B_\infty(f)$ if and only if there exists a sequence of points $(p_k)_{k\in\mathbb{N}} \subset \mathbb{C}^2$ such that $\|p_k\| \to \infty, \mathrm{grad}\ f(p_k) \to 0$ and $f(p_k) \to a$ as $k \to \infty$.

Upper bounds for $\#B_\infty(f)$ have been found in the 1990's by Lê V.T. and M. Oka [LO] in terms of Newton polyhedra at infinity. An estimation in terms of the degree d of f was given by Gwoździewicz and Płoski [GP]: if $\dim \mathrm{Sing}f \leq 0$ then $\#B_\infty(f) \leq \max\{1, d-3\}$. In the general case (dropping the condition $\dim \mathrm{Sing}f \leq 0$) we have $\#B_\infty(f) \leq d-1$, see e.g. [Je], [JK]. Recently Gwoździewicz [Gw] proved the following estimation of $\#B_\infty(f)$: if ν_0 denotes the number of branches at infinity of the fibre $f^{-1}(0)$, then the number of critical values at infinity other than 0 is at most ν_0. Here we refine and improve this statement by using a different method, in which results by Miyanishi [Mi1], [Mi] and Gurjar [Gu] play an important role.

For $a \in \mathbb{C}$, let us denote by ν_a the number of branches at infinity of the fiber $f^{-1}(a)$. This number is equal to ν_{gen} for all values $a \in \mathbb{C}$ except finitely many for which one may have either $\nu_a < \nu_{\mathrm{gen}}$ or $\nu_a > \nu_{\mathrm{gen}}$. Let $\nu_{\mathrm{min}} := \inf\{\nu_a \mid a \in \mathbb{C}\}$. Let us denote by b the number of *points at infinity of* f, i.e. $b := \#\overline{f^{-1}(a)} \cap L_\infty$, where L_∞ is the line at infinity $\mathbb{P}^n \setminus \mathbb{C}^n$.

Under these notations, our main result is the following:

Theorem 1.1. *Let $f : \mathbb{C}^2 \to \mathbb{C}$ be a polynomial function of degree d. Then:*

(a) $\#B_\infty(f) \leq \min\{\nu_{\mathrm{gen}}, \nu_{\mathrm{min}} + 1\}$.
(b) $\#\{a \in \mathbb{C} \mid \nu_a < \nu_{\mathrm{gen}}\} \leq \nu_{\mathrm{gen}} - b$.
(c) $\#\{a \in \mathbb{C} \mid \nu_a > \nu_{\mathrm{gen}}\} \leq \nu_{\mathrm{min}}$.

Date: January 28, 2014.
The first author was partially supported by Université Lille 1 and by the grant of NCN, 2014-2017.

In case $\nu_{\text{gen}} > \frac{d}{2}$*, we moreover have:*

(d) $\#B_\infty(f) \leq \min\{\nu_{\text{gen}} - 1, \nu_{\text{min}}\}$.
(e) $\#\{a \in \mathbb{C} \mid \nu_a > \nu_{\text{gen}}\} \leq \nu_{\text{min}} - 1$.

REMARK 1.2. Point (a) of Theorem 1.1 is equivalent to Gwoździewicz's [Gw, Theorem 2.1]. His result is a by-product of the local study of pencils of curves of Yomdin-Ephraim type. Our method is totally different and allows us to prove moreover several new issues, namely (b)–(e) of Theorem 1.1.

REMARK 1.3. As Gwoździewicz remarks, his inequality [Gw, Theorem 2.1] is "almost" sharp, i.e. not sharp by one. Our new inequality (d) improves by one the inequality (a) under the additional condition $\nu_{\text{gen}} > \frac{d}{2}$, thus yields the sharp upper bound, as shown by the example $f : \mathbb{C}^2 \to \mathbb{C}$, $f(x, y) = x + x^2 y$, where $d = \deg f = 3$, $\nu_{\text{min}} = \nu_{\text{gen}} = 2$, $b = 2$ and $B_\infty(f) = \{0\}$ with $\nu_0 = 3$.

The same example shows that our estimations (b) and (e) are also sharp.

Acknowledgements. The authors are grateful to Professor R.V. Gurjar from Tata Institute and Professor K. Palka from IMPAN for helpful discussions.

2. Proof of Theorem 1.1

We need here the important concept of *affine surfaces which contain a cylinder-like open subset* which was introduced by M. Miyanishi [Mi1]. Let us recall it together with some properties which we shall use.

Definition 2.1. [Mi] Let X be a normal affine surface. We say that X contains a *cylinder-like open subset* U, if there exists a smooth curve C such that $U \cong \mathbb{C} \times C$.

Let X be as in the above definition and let $\pi : U \to C$ be the projection. After [Mi, p.194], the projection π has a unique extension to a $\mathbb{C}$-fibration $\rho : X \to \bar{C}$, where $\bar{C}$ denotes the smooth completion of the curve C. We have the following important result of Gurjar and Miyanishi:

Theorem 2.2. [Mi1] [GM] [Gu] *Let X be a normal affine surface with a $\mathbb{C}$-fibration $f : X \to B$, where B is a smooth curve. Then:*

(a) *X has at most cyclic quotient singularities.*
(b) *Every fiber of f is a disjoint union of curves isomorphic to $\mathbb{C}$.*
(c) *A component of a fiber of f contains at most one singular point of X. If a component of a fiber occurs with multiplicity 1 in the scheme-theoretic fiber, then no singular point of X lies on this component.* □

Corollary 2.3. *Let X be a normal affine variety, which contains a cylinder-like open subset U. Then the set $X \setminus U$ is a disjoint union of curves isomorphic to $\mathbb{C}$. Moreover, every connected component l_i of this set contains at most one singular point of X.* □

Let $f : \mathbb{C}^2 \to \mathbb{C}$ be a polynomial function in fixed affine coordinates and denote by $\tilde{f}(x, y, z)$ the homogenization of f by a new variable z, namely $\tilde{f}(x, y, z) = f_d + z f_{d-1} + ... + z^d f_0$. Let $X := \{([x : y : z], t) \in \mathbb{P}^2 \times \mathbb{C} \mid \tilde{f}(x, y, z) = t z^d\}$ be the closure in $\mathbb{P}^2 \times \mathbb{C}$ of the graph $\Gamma := \text{graph}(f) \subset \mathbb{C}^2 \times \mathbb{C}$. Then X is a hypersurface and the points at infinity of f are precisely the set $\{a_1, ..., a_n\} := \{f_d = 0\} \subset L_\infty$, where $L_\infty := \{z = 0\}$ denotes the line at infinity. The projection $\pi : X \to \mathbb{C}$, $(x, t) \mapsto t$, is a proper extension of f.

Let X' be the normalization of X. Composing the normalisation map with π yields $\pi' : X' \to \mathbb{C}$, which is also a proper extension of f. For simplicity we shall denote it again by f in the following.

Lemma 2.4. *The set $X' \setminus \Gamma$ is a disjoint union of affine lines, $l_1, ..., l_r$, each line l_i is isomorphic to $\mathbb{C}$. On each line l_i there is at most one singular point of X'. Moreover, $b \leq r \leq \nu_{\min}$.*

Proof. Let us choose a line $l \subset \mathbb{P}^2$ such that $l \cap \{a_1, ..., a_n\} = \emptyset$. Let $X_1 := (\mathbb{P}^2 \setminus l) \times \mathbb{C} \cap X'$. The surface X_1 is affine and $X_1' \setminus \Gamma = \bigcup_{i=1}^r l_i$, where X_1' denotes the normalization of X_1. Since the surface X_1' contains a cylinder-like open subset $U \cong \mathbb{C}^2 \setminus l = \mathbb{C} \times \mathbb{C}^*$, the first part of our claim follows from Corollary 2.3. Next, the map f restricted to l_i is finite, hence surjective. This implies that every fiber of f has a branch at infinity which intersects l_i. In particular $r \leq \nu_{\min}$. The inequality $r \geq b$ is obvious. □

Denote by $f_i : l_i \cong \mathbb{C} \to \mathbb{C}$ the restriction of f to l_i. It can be identified with a one variable polynomial, the degree of which is equal to the number ν_i of branches of a generic fiber of f which intersect l_i. In particular $\sum_{i=1}^r \nu_i = \nu_{\text{gen}}$. The polynomial f_i of degree ν_i can have at most $\nu_i - 1$ singular points and their images by f_i are bifurcation values at infinity for f (this can be easily deduced from [Eh]). There might also be at most one singular point of X' which lies on l_i, and then its image by f is possibly a bifurcation value too. Summing up, we get that f can have at most ν_{gen} critical values at infinity, which shows one of the inequalities of point (a). Moreover, the inequality $\nu_a < \nu_{\text{gen}}$ is possible only if a is a critical value of some polynomial f_i. This means that $\#\{a \in \mathbb{C} \mid \nu_a < \nu_{\text{gen}}\} \leq \sum_{i=1}^r (\nu_i - 1) \leq \nu_{\text{gen}} - r \leq \nu_{\text{gen}} - b$. This proves (b).

Let us assume now $\nu_a = \nu_{\min}$. We have $\nu_a \geq \sum_{i=1}^r \#\{x \in l_i \mid f_i(x) = a\}$ since in every such point x there is at least one branch at infinity of the fiber $f^{-1}(a)$. Note that if $f_i(x) = a$ then $\operatorname{ord}_x(f_i - a) = \operatorname{ord}_x f_i' + 1$. Thus:

$$\#\{x \in l_i \mid f_i(x) = a\} = \sum_{x \in l_i, f_i(x) = a} [\operatorname{ord}_x(f_i - a) - \operatorname{ord}_x f_i'].$$

We have clearly the equality $\sum_{x \in l_i} \operatorname{ord}_x(f_i - a) = \nu_i$. Hence

$$\sum_{x \in l_i, f_i(x) = a} [\operatorname{ord}_x(f_i - a) - \operatorname{ord}_x f_i'] = \nu_i - \sum_{x \in l_i, f_i(x) = a} \operatorname{ord}_x f_i'.$$

Since $\sum_{x \in l_i} \operatorname{ord}_x f_i' = \nu_i - 1$ we have:

$$\nu_i - \sum_{x \in l_i, f_i(x) = a} \operatorname{ord}_x f_i' = 1 + \sum_{x \in l_i, f_i(x) \neq a} \operatorname{ord}_x f_i'.$$

Note that:

$$1 + \sum_{x \in l_i, f_i(x) \neq a} \operatorname{ord}_x f_i' \geq \#\{x \in l_i \mid f(x) \neq a, \text{ and either } f_i'(x) = 0 \text{ or } x \in \operatorname{Sing}(X')\}.$$

The number at the right side is greater or equal to the number of critical values at infinity of f different from $f^{-1}(a)$. Finally, taking the sum over all $i \in \{1, \ldots, r\}$ we get $\#B_\infty(f) \leq \nu_{\min} + 1$, which completes the proof of (a).

To prove (c), note that if the fiber $f^{-1}(a)$ does not contain singular points of X', then the intersection multiplicity $\overline{l_i} \cdot f^{-1}(a)$ is equal to $\nu_i = \deg f_i$, hence the fiber $f^{-1}(a)$ has at most ν_i branches on l_i. This implies $\nu_a \leq \nu_{\text{gen}}$. Hence $\#\{a \in \mathbb{C} \mid \nu_a > \nu_{\text{gen}}\} \leq r \leq \nu_{\min}$.

To prove (d) and (e) it is enough to show that if $\nu_{\text{gen}} > \frac{d}{2}$, then at least one line l_i does not contain singular points of X'. Let d_i be the smallest positive integer such that $d_i l_i$ is a Cartier divisor in X' (such a number exists because X' has only cyclic singularities). Since l_i is smooth, we have that $d_i = 1$ if and only if the line l_i does not contain any singular point of X', see Lemma 2.5 below. Now let Z be the closure of Γ in $\mathbb{P}^2 \times \mathbb{P}^1$ and let Z' denote its normalization. We have clearly the inclusion $X' \subset Z'$. Let $\Pi : Z' \to \mathbb{P}^2$ the first projection, where the second projection $Z' \to \mathbb{P}^1$ is an extension of f which we will denote for simplicity by f again. Let $f^{-1}(\infty) = S_1 \cup ... \cup S_k$ (where S_i are irreducible and taken with reduced structure). Recall that $L_\infty = \mathbb{P}^2 \setminus \mathbb{C}^2$ is the line at infinity. We have $\Pi^*(L_\infty) = \sum_{i=1}^k m_i S_i + \sum_{i=1}^r e_i \overline{l_i}$. Since $\Pi^*(L_\infty)$ is a Cartier divisor we have $e_i = n_i d_i$, where n_i is a positive integer.

Let us assume that any line l_i contains some singular point of X', i.e., that $d_i > 1$ for any i. Denoting by $F \subset \mathbb{P}^2$ the closure of a general fiber of $f = a$, since Π is a birational morphism, we have:

$$d = F \cdot L_\infty = \Pi^*(F) \cdot \Pi^*(L_\infty) = f^*(a) \cdot (\sum_{i=1}^k m_i S_i + \sum_{i=1}^r e_i \overline{l_i}).$$

Note that $\Pi^*(F) \cdot \sum_{i=1}^k m_i S_i = 0$ since $|f^*(a)| \cap |\sum_{i=1}^k m_i S_i| = |f^*(a)| \cap |f^*(\infty)| = \emptyset$. Moreover we have $\nu_i = f^*(a) \cdot \overline{l_i}$. Thus:

$$d = \sum_{i=1}^r n_i d_i \nu_i \geq \sum_{i=1}^r 2\nu_i = 2\nu_{\text{gen}}$$

and this ends our proof. □

Lemma 2.5. *Let X^n be an algebraic variety and let $Z^r \subset X^n$ be a subvariety which is a complete intersection in X^n. If a point $z \in Z^r$ is nonsingular on Z^r, then it is nonsingular on X^n.*

Proof. Let us recall that if X^n is an algebraic variety, then a point $z \in X^n$ is smooth, if and only if the local ring $\mathcal{O}_z(X)$ is regular. This is equivalent to the fact that $\dim_{\mathbb{C}} \mathfrak{m}/\mathfrak{m}^2 = \dim X^n = n$, where $\mathfrak{m}$ denotes the maximal ideal of $\mathcal{O}_z(X^n)$.

Let $I(Z^r) = \{f_1, ..., f_l\} \subset \mathbb{C}[X^n]$, where $l = n - r$, be the ideal of Z^r in the ring $\mathbb{C}[X^n]$. We have $\mathcal{O}_z(Z^r) = \mathcal{O}_z(X^n)/(f_1, ..., f_l)$. In particular if $\mathfrak{m}'$ denotes the maximal ideal of $\mathcal{O}_z(Z^r)$ and $\mathfrak{m}$ denotes the maximal ideal of $\mathcal{O}_z(X^n)$ then $\mathfrak{m}' = \mathfrak{m}/(f_1, ..., f_l)$. Let α_i denote a class of the polynomial f_i in $\mathfrak{m}/\mathfrak{m}^2$. Let us note that

$$\mathfrak{m}'/\mathfrak{m}'^2 = \mathfrak{m}/(\mathfrak{m}^2 + (\alpha_1, ..., \alpha_l)). \tag{2.1}$$

Since the point z is smooth on Z^r we have $\dim_{\mathbb{C}} \mathfrak{m}'/\mathfrak{m}'^2 = \dim Z^r = \dim X^n - l$. Take a basis $\beta_1, ..., \beta_{n-l}$ of the space $\mathfrak{m}'/\mathfrak{m}'^2$ and let $\overline{\beta_i} \in \mathfrak{m}/\mathfrak{m}^2$ correspond to β_i under the correspondence (2.1). Note that vectors $\overline{\beta_1}, ..., \overline{\beta_{n-l}}, \alpha_1, ..., \alpha_l$ generate the space $\mathfrak{m}/\mathfrak{m}^2$. This means that $\dim_{\mathbb{C}} \mathfrak{m}/\mathfrak{m}^2 \leq n - l + l = n = \dim X^n$. Since we also have the general inequality $\dim_{\mathbb{C}} \mathfrak{m}/\mathfrak{m}^2 \geq \dim X^n$, we obtain the equality $\dim_{\mathbb{C}} \mathfrak{m}/\mathfrak{m}^2 = \dim X^n$, which means that X^n is nonsingular at z. □

References

[CK] Chądzyński , T. Krasiński, *The gradient of a polynomial at infinity*, Kodai Math.J. 26 (2003), 317-339.

[Du] A.H. Durfee, *Five definitions of critical point at infinity*, Singularities (Oberwolfach, 1996), 345-360, Progr. Math., 162, Birkhuser, Basel, 1998.

[Eh] C. Ehresmann, *Sur lespaces fibrés différetiables*, C.R. Acad. Sci. Paris 224 (1947), 1611-1612.
[HL] Hà H.V., Lê D.T., *Sur la topologie des polynômes complexes*, Acta Math. Vietnam. 9 (1984), no. 1, 21-32.
[Gu] R. V. Gurjar, *Remarks on the Topology of Surface singularities and Applications*, Proc. of Indo-French conference on Geometry, NBHM, (1993), 95-102.
[GM] R.V. Gurjar, M. Miyanishi, *Automorphisms of Affine Surfaces with $\mathbb{A}^1$-Fibrations*, Michigan Math. J. 53 (2005), 33-55.
[GP] J. Gwoździewicz, A. Płoski, *Formulae for the singularities at ifinity of plane algebraic curves*, Univ. Iagell. Acta Math. XXXIX, (2001), 109-133.
[Gw] J. Gwoździewicz, *Ephraim's pencils*, Int. Math. Res. Notices 15 (2013), 3371-3385.
[Je] Z. Jelonek, *On bifurcation points of a complex polynomial*, Proc. AMS. 131, (2003), 1361 - 1367.
[JK] Z. Jelonek, K. Kurdyka, *On asymptotic critical values of a complex polynomial*, J. Reine Angew. Math. 565, (2003), 1-11.
[LO] Lê V.T., M. Oka, *Estimation of the Number of the Critical Values at Infinity of a Polynomial Function*, Publ. RIMS. Kyoto Univ. 31 (1995), 577-598.
[Mi] M. Miyanishi, *Open algebraic surfaces*, CRM Monograph Series, 12. American Mathematical Society, Providence, RI, 2001. viii+259 pp.
[Mi1] M. Miyanishi, *Singularities of normal affine surfaces containing cylinderlike open sets*. J. Algebra 68 (1981), 268-275.
[Su] M. Suzuki, *Propriétés topologiques des polynômes de deux variables complexes, et automorphismes algébriques de l'espace $\mathbb{C}^2$*, J. Math. Soc. Japan 26 (1974), 241-257.
[T] R. Thom, *Ensembles et morphismes stratifiés*, Bull. Amer. Math. Soc. **75** (1969), 249-312.
[Ti1] M. Tibăr, *Regularity at infinity of real and complex polynomial functions*, Singularity theory (Liverpool, 1996) 249-264, London Math. Soc. Lecture Note Ser., 263, Cambridge Univ. Press, Cambridge, 1999.
[Ti2] M. Tibăr, Polynomials and vanishing cycles, Cambridge Tracts in Mathematics, 170. Cambridge University Press, Cambridge, 2007. xii+253 pp.

Instytut Matematyczny, Polska Akademia Nauk, Śniadeckich 8, 00-956 Warszawa, Poland.
E-mail address: najelone@cyf-kr.edu.pl

Mathématiques, UMR 8524 CNRS, Université Lille 1, 59655 Villeneuve d'Ascq, France.
E-mail address: tibar@math.univ-lille1.fr